

590 1136

CYC 933 30

83467 84427

Reports from the Environmental Archaeology Unit, York

Technical Report: Plant and invertebrate remains from excavations associated with renovations at All Saints Church, Pavement, York (site code 95.47)

by Allan Hall, Harry Kenward and John Carrott

Report 98/30

*Environmental Archaeology Unit
University of York
Heslington
York YO10 5DD*

*(01904) 433846/51
Fax: 433850*

Reports from the Environmental Archaeology Unit, York

Technical Report: Plant and invertebrate remains from excavations associated with renovations at All Saints Church, Pavement, York (site code 95.47)

by Allan Hall, Harry Kenward and John Carrott

Report 98/30

*Environmental Archaeology Unit
University of York
Heslington
York YO10 5DD*

*(01904) 433846/51
Fax: 433850*

This report is part of the series *Reports from the Environmental Archaeology Unit, York*

To avoid confusion as to the nature of the various kinds of reports produced by the EAU, their typical level and content is outlined here.

An *evaluation report* presents the results of examination of hand-collected and sieved material, and of examination and processing of sediment samples, from an evaluation excavation. (An evaluation excavation is defined as one carried out to determine the archaeological importance and potential of deposits in advance of development but not intended to produce a definitive account of the deposits.) The environmental archaeology component of evaluations is carried out to standardised and very limited specifications and budgets and the data obtained are generally at least somewhat subjective. However, since it is the experience of EAU staff that few evaluations are followed by any more systematic excavation and research, an attempt is made to produce a report which contains sufficient information to be of at least some value in synthesis. Where possible, material of particular importance is examined and reported more fully than covered by the costing, using overhead funding. Recommendations are made as to the desirability of retaining material for possible future research. No attempt can be made in an evaluation report to do more than give the most general information relating the site to others; the archaeological information available when the evaluation report is written by EAU is usually very limited.

An *assessment report* presents the results of the first stage of post-excavation work, often carried out under English Heritage's 'MAP2' scheme. Archaeological information is generally available at a basic descriptive level (perhaps only context type), with preliminary phasing and dating. Practical work is carried out on material selected in conjunction with the post-excavation team, and material is examined (a) to determine its potential for interpretation, and thus (by extrapolation) that of the environmental samples as a whole; and (b) to permit the scale and cost of the main phase of post-excavation work to be estimated. Practical work is carried out as rapidly as is compatible with realising these aims whilst maintaining an acceptable level of accuracy. It is assumed that there is a reasonable likelihood of any significant material being examined in the main phase of post-excavation work, so the assessment report cannot be seen as being at all

a definitive account of the environmental archaeology of the site.

A *technical report* represents the basic reporting of the environmental archaeology evidence, with a greater or lesser amount of analytical, comparative and synthetic material. Many technical reports are written so as to allow extraction of sections of text for a publication report, but this should not be done without reference to the EAU. Some technical reports do no more than place evidence on record, and they may concern deposits with very limited potential.

Data archives put datasets on record, with, in most cases, minimal accompanying text. *Notes* report methodological and other occasional observations and may have almost any format.

Publication texts are not normally included in this series except as *drafts* or *preprints*. These are copies of material submitted or accepted for publication respectively. It should not be assumed that the first will be published in the form presented, or at all, and some illustrative material, particularly photographs, may be poorly reproduced or omitted from both.

To the client:

This report represents in good faith the product as defined implicitly or explicitly in the contract or other agreement between the client and the EAU. Any errors or deficiencies will be rectified as soon as reasonably possible. Revisions necessitated through late changes to phasing or other archaeological information may result in an additional charge. Unless otherwise stated, *minor* reworking of technical report material for publication is covered by the initial contract.

How to obtain copies of Reports from the EAU, York

Much of the material in this series is commissioned by clients on behalf of developers, and may be confidential. In general, however, copies are obtainable at cost (photocopying, labour and postage) from the EAU. A full list of reports originating from the EAU, and of publications by EAU staff, is available on request.

Reports from the Environmental Archaeology Unit, York 98/30, 20 pp.

**Technical Report: Plant and invertebrate remains
from excavations associated with renovations
at All Saints Church, Pavement, York (site code 95.47)**

by Allan Hall, Harry Kenward and John Carrott

Summary

Samples collected during excavation and piling consequent upon renovation of All Saints Church, Pavement, York, were submitted for assessment of their bioarchaeological value. Three samples of deposits thought to date to the time before the construction of the earliest known church and probably to the Anglo-Scandinavian period yielded plant and invertebrate assemblages of considerable interpretative value and were examined subsequently in more detail. They provide bioarchaeological evidence closely similar to that from deposits of Anglo-Scandinavian date from the south side of Coppergate, having good preservation of a rich variety of plant remains, with abundant wood and sometimes also bark, and the presence of remains of at least four plants interpreted as having been used in dyeing and mordanting of textiles. The insect assemblages were of limited size and variable preservation but probably accumulated through deposition of various materials in the open air.

The remaining samples, from grave/coffin fills from within the church, were almost barren except for the presence of some human bone in some instances.

Keywords: ALL SAINTS CHURCH; YORK; ENGLAND (NE); ANGLO-SCANDINAVIAN; MEDIEVAL; ?POST-MEDIEVAL; GRAVE FILLS; OCCUPATION DEPOSITS; MICROFOSSILS; PLANT REMAINS; INSECT REMAINS

Prepared by:

Environmental Archaeology Unit
University of York
Heslington
York YO10 5DD

For:

MAP Archaeological Consultancy Ltd.
39 Greengate
MALTON
N. Yorks YO17 0EL

24th September 1998

Technical Report: Plant and invertebrate remains from excavations associated with renovations at All Saints Church, Pavement, York (site code 95.47)

Introduction and methods

Excavations were undertaken by MAP Archaeological Consultancy Ltd. during 1995 within and to the south of All Saints Church, Pavement, York, during renovation and underpinning operations carried out with financial support from English Heritage. A series of small samples from test boreholes was submitted to the Environmental Archaeology Unit, University of York, in 1995 for assessment of their bioarchaeological value and an informal report compiled (included as an appendix to the assessment report, see below). A number of deposits from grave or coffin fills (from trenches) or from deposits apparently pre-dating the earliest church on the site (from pile-borings) were sampled by means of 'GBAs' (*sensu* Dobney *et al.* 1992) and these were assessed for their bioarchaeological potential in 1996 (Carrott *et al.* 1996). Latterly, further material from three of these samples (selected on the basis of the results of the assessment exercise) has been examined in more detail and it is the combined results of all these investigations that are presented in this report.

The GBAs have been examined by means of 'test' subsamples (Kenward *et al.* 1986), using either a 'flot' from paraffin flotation or a 'washover' to provide fractions for assessment of insect remains; in addition, one tubful of one of the samples (21) was 'bulk-sieved' to 0.5 mm. Two samples were described but not subjected to further analysis since they appeared very unlikely to contain interpretatively useful biological remains and it was necessary to constrain costs at the assessment stage.

Plant remains and other components were examined in the residues, flots and washovers; flots and washovers were checked for insect remains (which were also, to a limited extent,

recovered from residues). 'Squashes' (*sensu* Dainton 1992) were made on selected samples to check for the presence of microfossils, especially parasite eggs.

Results

The results of the various analyses of GBA samples are presented in context number order, with comments from the excavator in square brackets. The work undertaken is indicated by the paragraph headings 'Assessment' and 'Assessment and analysis'. Data are presented in Tables 1-6.

Context 2054 [fill of grave cut around articulated skeleton 2055; ?post-medieval; routine sampling of burial]

Sample 23: Mid grey-brown, unconsolidated sand with stones 2-60 mm present, together with some brick/tile fragments. A 1 kg 'test' subsample was processed and the excess material bulk-sieved to 1 mm.

Assessment: There was a large residue of sand, with some brick/tile and mortar and a trace of charcoal to 10 mm, and a little (?human) bone. The tiny washover (a few cubic centimetres at most) included traces of charcoal (<5 mm), bone and an elderberry (*Sambucus nigra*) seed, together with a few traces of invertebrate cuticle. The squash consisted of inorganic particles with a trace of organic detritus. No microfossils were observed.

Context 4133 [grave fill of complete articulated ?medieval skeleton 4138; routine sampling of fill]

Sample 20: Mid golden-grey-brown, unconsolidated, stony sand with stones 2-60+ mm and mortar/plaster present. A 1 kg 'test' subsample was processed.

Assessment: There was a large residue of sand and mortar with traces of gravel, brick/tile and charcoal (<10 mm); the very small flot contained a few fragments of very decayed wood (<5 mm) and charcoal (<10 mm) and a single sedge (*Carex*) nutlet. There were no more than traces of invertebrate remains. The squash consisted mostly of inorganic particles, with a little organic detritus. Some silica bodies resembling phytoliths were noted, but the identification was not certain.

Context 4139 [clean sand bedding at base of grave on which nearly complete skeleton 4137 rested; only burial of this kind; routine sampling of grave fill]

Sample 22: Mid golden-brown, unconsolidated coarse sand with stones 6-60+ mm present, together with some large chunks of mortar/plaster. Not investigated further.

Context 4143 [fill of coffin around complete articulated ?medieval skeleton 4146; routine sampling of burial]

Sample 21: Mid grey, crumbly, unconsolidated sand with mortar/plaster and human bone present (a second tub: mid golden-brown, unconsolidated sand with stones 20-60 mm, mortar/plaster, brick/tile, coal and large mammal bone present). A 1 kg 'test' subsample was processed.

Assessment: The large residue was of sand and mortar with some ?human bone, a trace of charcoal (<10 mm) and brick/tile; the tiny washover (a few cubic centimetres only) contained some charcoal and very decayed wood (<5 mm). There were a few fragments of a ?modern millipede but no other invertebrate remains were observed. The squash consisted primarily of inorganic particles, although there was some organic material. A few phytoliths were noted.

Context 4164 [material cut by earliest structure encountered; quite unlike other deposits recorded from within church, ?pre-church and Anglo-Scandinavian in date]

Sample 24: Dark grey-brown, crumbly, humic slightly sandy silt with patches of matted

herbaceous detritus locally, iron-rich concretions (perhaps an iron object), charcoal, large mammal bone and oyster shell fragments. A 1 kg 'test' subsample was processed.

Assessment: The small residue (of about 0.2 litres) consisted of about half its volume of organic material, mainly charcoal to 15 mm, with some very decayed wood to 15 mm; there was some bone and a few rather poorly preserved weed seeds. There was also a single charred fragment of a woody stem which may have been dyer's greenweed (*Genista tinctoria*), a plant recorded from some of the other pre-church deposits (see below). The tiny flot yielded a few seeds and invertebrate remains of no interpretative value. The lack of 'waterlogged' plant and insect remains is rather surprising given the presence of 'matted herbaceous detritus' observed in the raw sediment.

Context 4171 [thin lens beneath Context 4164 (see above); material cut by earliest structure encountered; ?pre-church and Anglo-Scandinavian in date; ?ash]

Sample 25: Light grey ash with darker patches of charcoal. Not investigated further.

Context 4172 [slightly laminated organic deposit sealed by Context 4171 (see above); material cut by earliest structure encountered; quite unlike other deposits recorded from within church, ?pre-church and Anglo-Scandinavian in date]

Sample 26: Dark grey (speckled light grey at mm scale), crumbly, soft, working just plastic, humic sandy silt with inclusions of herbaceous detritus, a slab of flaggy sandstone (with pale grey ash or mortar associated with it), charcoal, well decayed wood, large mammal bone and oyster shell fragments. Subsamples of 1 and 5.8 kg were processed.

Assessment and analysis: The residues of the combined subsamples consisted of about 2.5 litres of material of which about 40% by volume was organic—mainly decayed wood (to 30 mm) with some charcoal to 20 mm—and another 40% oyster shell with some bone. There was also some bark,

and amongst the wood fragments were chips from working of wood.

There was good evidence for a suite of plants associated with textile dyeing and which were common in Anglo-Scandinavian deposits at 16-22 Copper-gate, a little more than 20 metres to the south of the present site: moderate numbers of twig fragments of dyer's greenweed, plus traces of stem fragments of the clubmoss *Diphasium complanatum* and root fragments of madder (*Rubia tinctorum*). The tiny flot yielded several *Diphasium* fragments and some well preserved achenes of hop (*Humulus lupulus*). Other plants, mostly present in very low concentrations, were a mixture of weeds, wetland and food plants.

Insect remains were rather rare in the flot from the 1 kg assessment subsample, but it was considered that an interpretable fauna would be recovered from a much larger subsample. In the event, 5.8 kg processed for analysis produced only about twice as many insect fragments as were recorded from the initial subsample, probably at least in part as the result of presence of a large volume of oyster shell and wood. Although this group would not be out of place amongst the assemblages from Anglo-Scandinavian Coppergate (Kenward and Hall 1995), it had no clear character. It is notable that, although the material from the first subsample was described as showing 'good to moderate preservation', the remains in the second were 'rather decayed and very fragmented', probably suggesting that the sample was heterogeneous (decay during storage is considered unlikely in this case).

Other material, presumably from occupation, recorded in the residue included charcoal, apple (*Malus*) endocarp (core fragments), mammal bone, mortar, oyster shell fragments, and pottery.

Context 5012 [from pile-hole 5, 12.25 m OD; organic silt from under first church, ?Anglo-Scandinavian]

Sample 28: Dark grey, crumbly (working just plastic), humic, slightly sandy clay silt with mortar/plaster and large mammal bone present. Two separate subsamples of 1 kg were processed.

Assessment and analysis: There was a moderately large combined residue (of about 0.7 litres) of which about two-thirds by volume was organic: charcoal (to 20 mm), bark (to 50 mm) and decayed wood fragments (to 20 mm); the wood included some ?modern pale conifer wood. Small numbers of generally well preserved seeds were present, mostly probably from weed taxa; however, there were also some achenes of hop, a capsule fragment of flax/linseed (*Linum usitatissimum*), and fragments of dyer's greenweed and clubmoss stem and madder root. There was a single charred ?wheat (*Triticum*) grain and one well-preserved oat (*Avena*) grain. A quite long list of other plant taxa was recorded (Table 2), most of them weeds, though with some heather, food plants (hazelnut, apple endocarp) and a moderately wide range of mosses, most of which were types widely recorded at 16-22 Coppergate. The residue also contained some charcoal fragments, some (unusually) bearing a blue residue (presumably vivianite); the rest of the residue comprised sand with some mortar, brick/tile, mammal bone, gravel and pottery. The tiny flots contained a few rather poorly preserved seeds, including more hop achenes, and further rare fragments of *Diphasium*.

There were moderate quantities of insect remains in the flot from the first subsample, both adults and immatures. Most were of species associated with decomposing matter, and this component was subjectively very similar to many groups recorded from Anglo-Scandinavian deposits at 16-22 Coppergate. There were also two *Helophorus* water beetles, doubtless 'background fauna' elements. Of the more abundant taxa, *Acritus nigricornis* (4 individuals) is associated with somewhat foul, decaying matter, and *Anotylus nitidulus* (3) probably exploited similar conditions. *Xylodromus concinnus* (4) was probably associated with rather drier material. It is regarded as typical of 'house fauna' assemblages and the presence of such a component was supported by single individuals of a few other taxa. The second subsample gave remains showing good preservation. The assemblage was not very similar to that from the first subsample, the most abundant beetles being *Trox scaber* (3), *Carpelimus fuliginosus* (also 3), and *C. ?bilineatus* (4). There was probably somewhat foul decaying matter at

the point of deposition but no clear interpretation can be offered. Again, there were house fauna elements, including a single human flea, *Pulex irritans*.

Context 5015 [from pile-hole 6; 12.25 m OD; dark organic deposit under foundations of first church, ?Anglo-Scandinavian date]

Sample 27: Dark grey, crumbly, humic, slightly sandy silt with patches of fine herbaceous detritus locally, and charcoal, wood and large mammal bone all present. Subsamples of 1 kg and 0.8 kg were processed.

Assessment and analysis: Of the moderately large residue (of about 1 litre), about three-quarters was decayed wood, including well-preserved wood chips (to 35 mm), and two large fragments of vertebrate (a butchered cow bone and a red deer antler fragment) which made up about 10-15% of the volume of the smaller subsample; other debris included *Diphysium complanatum* and *Genista tinctoria* stem fragments and *Rubia* root. There was only a low concentration of rather poorly preserved seeds, though a quite wide range of plant taxa was present, including weeds, grassland forms (perhaps from hay and/or stable manure), food plants and wetland taxa. No one group predominated. Overall the assemblage was reminiscent of some of the floor deposits from 16-22 Coppergate. The remainder of the residue comprised sand and gravel with a little bone, pot and mortar.

The two squashes carried out showed approximately equal quantities of inorganic and organic particles; there were several 'grass-type' phytoliths and diatoms and a few fungal spores.

The tiny flots (a few cubic centimetres at most) from the first subsample was mainly of fine herbaceous detritus. Moderately well preserved invertebrate remains were present in modest numbers and, although the species composition was different from that seen in Sample 28, they, too, constituted an assemblage typical of Anglo-Scandinavian deposits. Subjectively there was a 'house fauna' component, with which a puparium of the sheep ked *Melophagus ovinus* should be

included, doubtless having originated in wool cleaning. Slightly foul conditions were indicated. A single honey bee (*Apis mellifera*) first hind tarsal segment was also noted.

The second subsample produced only rather few insect remains, the fauna having some subjective differences from that of the first in that there were hints of rather fouler material. There were traces of house fauna, including a human flea.

Context 5116 [from pile-hole 36, material of a kind not encountered in other holes or in excavations, ?pre-church, ?domestic]

Sample 29: Mid grey-brown, soft silty clay sand with stones 6-60 mm, ?human bone, and a lump of tufa making up about one-third of the sample. A 1 kg 'test' subsample was processed.

Assessment: The moderately large residue (about 0.5 litres) was essentially of sand with some gravel, mortar, brick/tile, ?human bone, charcoal (<10 mm). Some of the tufa was dissolved in dilute hydrochloric acid to check for the presence of identifiable remains trapped within its matrix but it was not productive. The small washover (a few cubic centimetres only) was of charcoal (to 5 mm) and very decayed wood (also to 5 mm), with a few rather poorly preserved seeds of no interpretative value. There were no more than traces of invertebrates.

Tufa of the kind observed from this pile-hole was recorded as 'spot' finds on several occasions during excavation of Anglo-Scandinavian deposits at 16-22 Coppergate (Kenward and Hall 1995, 719, 750). Its origin is unknown, though formation within the built-up town is suspected, perhaps in association with limestone buildings.

Discussion

Three of the samples examined during assessment (from Contexts 4172, 5012 and 5105) were considered to have potential for worthwhile bioarchaeological analysis and were selected for further investigation via additional subsamples. For the plant remains, the additional material

provided confirmation of the nature of the assemblages and, inherently also reinforced the belief that the material was of Anglo-Scandinavian date: the assemblages contained a diversity of taxa including traces of plants likely to have been used in textile dyeing and mordanting (of the kinds discussed by Kenward and Hall 1995). Wood (including chips from woodworking) and bark were also important, as in many of the Coppergate samples.

The results of the insect analyses also served to underline the typical 'Anglo-Scandinavian' character of the biota, but unfortunately in two cases the quantity of material remaining for processing was small and in the third the large additional subsample examined produced a disproportionately small number of remains by comparison with the assessment subsample. The insects thus gave only limited information about local ecology; it may be stated that none of the groups showed a strong character and that they probably represent external accumulations. To an extent, this is supported by the weak subjective similarity of the fauna from Sample 28 to some assemblages recorded at the nearby site at 5-7 Coppergate (Hall *et al.* 1983) and regarded as reasonably certainly having formed on a damp surface in the open air. The presence of human fleas in two of the samples is by no means incompatible with this; fleas seem to have been deposited in small numbers fairly widely on sites of this period and there was a range of 'house fauna' suggesting scattering or dumping of floor material. Similarly, a single honey bee does not provide evidence of bee-keeping, but is consistent with the numerous records of honey bees a few metres away at 16-22 Coppergate.

The grave/coffin fills examined were not productive of plant and invertebrate remains although there was some human bone.

Archive

All paper and electronic archive material pertaining to the work described here is stored, together with the remaining unprocessed sediment and any residues, flots or washovers, at the

Environmental Archaeology Unit, University of York.

Acknowledgments

The authors are grateful to Michael Issitt and Frances Large for undertaking practical work on these samples and to MAP Archaeological Consultancy Ltd. for providing material and for information about the archaeological contexts from which the samples came.

References

- Carrott, J., Hall, A., Issitt, M., Kenward, H. and Large, F. (1996). Assessment of plant and invertebrate remains from excavations associated with renovations at All Saints Church, Pavement, York (site code 95.47). *Reports from the Environmental Archaeology Unit, York* 96/47, 8 pp.
- Dainton, M. (1992). A quick, semi-quantitative method for recording nematode gut parasite eggs from archaeological deposits. *Circaea, the Journal of the Association for Environmental Archaeology* 9, 58-63.
- Dobney, K., Hall, A. R., Kenward, H. K. and Milles, A. (1992). A working classification of sample types for environmental archaeology. *Circaea, the Journal of the Association for Environmental Archaeology* 9 (for 1991), 24-6.
- Hall, A. R., Kenward, H. K., Williams, D. and Greig, J. R. A. (1983). Environment and living conditions at two Anglo-Scandinavian sites. *The Archaeology of York* 14 (4), 157-240 plus Plate I and Fiche 1. London: Council for British Archaeology.
- Kenward, H. K., Engleman, C., Robertson, A., and Large, F. (1986). Rapid scanning of urban archaeological deposits for insect remains. *Circaea* 3 (for 1985), 163-72.
- Kenward, H. K. and Hall, A. R. (1995). Biological evidence from Anglo-Scandinavian deposits at 16-22 Coppergate. *The Archaeology of York* 14 (7), 435-797 + xxii + loose figures. York: Council for British Archaeology.
- Kloet, G. S. and Hincks, W. D. (1964-77). *A check list of British Insects*. (2nd ed.) London: Royal Entomological Society.

Smith, A. J. E. (1978). *The moss flora of Britain and Ireland*. Cambridge: University Press.

Tutin, T. G. *et al.* (1964-80). *Flora Europaea 1-5*. Cambridge: University Press.

Table 1. Complete list of plant taxa recorded from All Saints Church, Pavement, York, in taxonomic order; nomenclature follows Tutin et al. (1964-90) for vascular plants and Smith (1978) for mosses .

<i>Diphasium complanatum</i> (L.) Rothm.	complanate clubmoss	shoot fragment(s)
<i>Betula</i> sp(p).	birch	fruit(s)
<i>Corylus avellana</i> L.	hazel	charred nut(s) and/or nutshell fragment(s)
<i>Corylus avellana</i> L.	hazel	nut(s) and/or nutshell fragment(s)
<i>Humulus lupulus</i> L.	hop	achene(s)
<i>Urtica dioica</i> L.	stinging nettle	achene(s)
<i>U. urens</i> L.	annual nettle	achene(s)
<i>Polygonum aviculare</i> agg.	knotgrass	fruit(s)
<i>P. hydropiper</i> L.	water-pepper	fruit(s)
<i>P. persicaria</i> L.	persicaria/red shank	fruit(s)
<i>P. lapathifolium</i> L.	pale persicaria	fruit(s)
<i>Bilderdykia convolvulus</i> (L.) Dumort.	black bindweed	fruit(s)
<i>Rumex acetosella</i> agg.	sheep's sorrel	fruit(s)
<i>Rumex</i> sp(p).	docks	fruit(s)
<i>Chenopodium album</i> L.	fat-hen	seed(s)
<i>Atriplex</i> sp(p).	oraches	seed(s)
<i>Stellaria media</i> (L.) Vill.	chickweed	seed(s)
<i>Spergula arvensis</i> L.	corn spurrey	seed(s)
<i>Agrostemma githago</i> L.	corncockle	seed(s), seed fragment(s)
<i>Silene</i> cf. <i>alba</i> (Miller) Krause in Sturm	?white campion	seed(s)
<i>Ranunculus</i> Section <i>Ranunculus</i>	meadow/creeping/bulbous buttercup	achene(s)
<i>R. sardous</i> Crantz	hairy buttercup	achene(s)
<i>R. flammula</i> L.	lesser spearwort	achene(s)
<i>Thlaspi arvense</i> L.	field penny-cress	seed(s)
<i>Coronopus squamatus</i> (Forskål) Ascherson	swine-cress	fruit(s)
cf. <i>C. squamatus</i>		seed(s)
<i>Brassica rapa</i> L.	'turnip'	seed(s)
<i>Brassica</i> sp(p).	cabbages, etc.	seed(s)
<i>Raphanus raphanistrum</i> L.	wild radish	pod segments and/or fragment(s)
<i>Reseda luteola</i> L.	weld/dyer's rocket	seed(s)
<i>R. lutea</i> L.	wild mignonette	seed(s)
<i>Rubus fruticosus</i> agg.	blackberry/bramble	seed(s)
<i>Potentilla</i> cf. <i>erecta</i> (L.) Räuschel	?tormentil	achene(s)
<i>Malus sylvestris</i> Miller	crab apple	endocarp
<i>Prunus spinosa</i> L.	sloe	fruitstone(s)
Leguminosae	pea family	pod(s) and/or pod fragment(s)
<i>Genista tinctoria</i> L.	dyer's greenweed	stem fragment(s), twig epidermis fragment(s)
cf. <i>G. tinctoria</i>		charred stem fragment(s), pod fragment(s)
<i>Vicia faba</i> L.	field bean	charred seed(s)
<i>Linum usitatissimum</i> L.	cultivated flax	capsule fragment(s)
<i>Aethusa cynapium</i> L.	fool's parsley	mericarp(s)
<i>Calluna vulgaris</i> (L.) Hull	heather, ling	bud(s)
cf. <i>C. vulgaris</i>		charred root and/or twig fragment(s)
<i>Galium</i> sp(p).	bedstraws, etc.	fruit(s)
<i>Rubia tinctorum</i> L.	dyer's madder	root fragment(s)
<i>Marrubium vulgare</i> L.	white horehound	nutlet(s)
<i>Galeopsis</i> Subgenus <i>Galeopsis</i>	hemp-nettles	nutlet(s)
<i>Prunella vulgaris</i> L.	selfheal	nutlet(s)
<i>Lycopus europaeus</i> L.	gipsywort	nutlet(s)
<i>Sambucus nigra</i> L.	elder	seed fragment(s), seed(s)
<i>Carduus/Cirsium</i> sp(p).	thistles	achene(s)

<i>Centaurea</i> sp(p).	knapweeds, etc.	achene(s)
<i>Sonchus asper</i> (L.) Hill	prickly sow-thistle	achene(s)
<i>Lapsana communis</i> L.	nipplewort	achene(s)
<i>Juncus bufonius</i> L.	toad rush	seed(s)
Gramineae	grasses	charred and waterlogged caryopsis/es
Cerealia indet.	cereals	charred caryopsis/es
<i>Triticum</i> sp(p).	wheats	charred caryopsis/es
<i>Hordeum</i> sp(p).	barley	charred caryopsis/es
<i>Avena</i> sp(p).	oats	charred caryopsis/es
<i>Scirpus lacustris</i> sensu lato	bulrush	nutlet(s)
<i>Eleocharis palustris</i> sensu lato	common spike-rush	nutlet(s)
<i>Carex</i> sp(p).	sedges	nutlet(s)

Mosses (all leaf/leaves and/or shoot fragment(s))

<i>Sphagnum</i> sp(p).
<i>Leucodon sciuroides</i> (Hedw.) Schwaegr.
<i>Neckera crispa</i> Hedw.
<i>N. complanata</i> (Hedw.) Hüb.
<i>Homalia trichomanoides</i> (Hedw.) Br. Eur.
<i>Thuidium</i> cf. <i>tamariscinum</i> (Hedw.) Br. Eur.
<i>Scorpidium scorpioides</i> (Hedw.) Limpr.
<i>Calliergon giganteum</i> (Schimp.) Kindb.
<i>C.</i> cf. <i>giganteum</i> (Schimp.) Kindb.
<i>Homalothecium sericeum</i> (Hedw.) Br. Eur. / <i>H. lutescens</i> (Hedw.) Robins.
<i>Brachythecium</i> sp(p).
<i>Hypnum</i> cf. <i>cupressiforme</i> Hedw.
<i>Rhytidiadelphus</i> sp(p).

Table 2. Plant remains and other components of the GBA test subsamples from All Saints Church, Pavement, York. Lists are presented in context order and for each subsample in descending order by 'amount' (on a four-point scale of abundance) and alphabetical order of component/taxon. For a complete list of plant taxa and an explanation of the abbreviated forms for parts recorded, see Table 1. Subsample weights are 1 kg unless otherwise indicated. Dim.—dimension; dec.—decayed; max.—maximum.

Context 2054 , Sample 23/T		?imbrex	1	max dim. 150 mm
sand	3	Aethusa cynapium	1	
brick/tile	2	Agrostemma githago (sf)	1	
mortar	2	Bilderdykia convolvulus	1	
bone fgts	1	bone fgts	1	max. size 100 mm
charcoal	1	Brassica rapa	1	
Sambucus nigra	1	Brassica sp(p).	1	
		Carex sp(p).	1	
		cf. Genista tinctoria (pod fgts)	1	max. size 15 mm
		Corylus avellana	1	
		Diphysium complanatum	1	
		earthworm egg caps	1	
		Eleocharis palustris sl	1	
		fish bone	1	
		fish scale	1	
		fly puparia	1	
		Fraxinus (charcoal)	1	
		Genista tinctoria (tef)	1	
		Gramineae (ch)	1	
		Homalia trichomanoides	1	
		Humulus lupulus	1	
		Juncus bufonius	1	
		Malus sylvestris (endo)	1	
		mammal bone	1	
		mortar	1	
		Polygonum aviculare agg.	1	
		Polygonum persicaria	1	
		Potentilla cf. erecta	1	
		pottery	1	
		Prunus spinosa	1	
		Rubia tinctorum	1	
		Rumex sp(p).	1	
		Sambucus nigra (sf)	1	
		Scirpus lacustris sl	1	
		wood chips	1	
Context 4133, Sample 20/T				
sand	4			
mortar	3			
brick/tile	1			
Carex sp(p).	1			
charcoal	1			max. size 10 mm
gravel	1			
Context 4143, Sample 21/T				
sand	4			
mortar	3			
?human bone	1			
brick/tile	1			
charcoal	1			max. size 10 mm
wood fgts	1			very decayed fragments
Context 4164, Sample 24/T				
charcoal	2			max. size 15 mm
Chenopodium album	2			
sand	2			
?human bone	1			
brick/tile	1			
cf. Genista tinctoria (ch st fgts)	1			
Eleocharis palustris sl	1			
fish bone	1			
mortar	1			
Sambucus nigra	1			
Stellaria media	1			
wood fgts	1			very decayed fragments
Context 4172, Sample 26/T (6.8 kg)				
charcoal	3			max. size 20 mm
oyster shell fgts	3			max. size 80 mm
wood fgts	3			max. size 30 mm
Atriplex sp(p).	2			
Chenopodium album	2			
Genista tinctoria (st fgts)	2			
sand	2			
Context 5012, Sample 28/T				
bark fgts	2			max. size 50 mm
charcoal	2			max. size 20 mm
sand	2			
Urtica dioica	2			
wood fgts	2			max. size 20 mm
Agrostemma githago	1			
Agrostemma githago (sf)	1			
Atriplex sp(p).	1			
Avena sp(p).	1			
beetles	1			
Bilderdykia convolvulus	1			
bone fgts	1			max. size 35 mm
Brassica sp(p).	1			
brick/tile	1			max. size 10 mm
Calliergon cf. giganteum	1			
Calluna vulgaris (b)	1			
Carex sp(p).	1			

Centaurea sp(p).	1		antler fgts	1	max. size 70 mm
cf. Calluna vulgaris			Atriplex sp(p).	1	
(ch rt-tw fgts)	1		bark fgts	1	max. size 20 mm
cf. Coronopus squamatus	1		beetles	1	
Chenopodium album	1		Betula sp(p).	1	
Coronopus squamatus (fr)	1		bone fgts	1	max dim. 110 mm
Corylus avellana	1		Brachythecium sp(p).	1	
Corylus avellana (ch)	1		Brassica sp(p).	1	
Diphasium complanatum	1		brick/tile	1	max. size 20 mm
earthworm egg caps	1		burnt bone fgts	1	max. size 10 mm
Eleocharis palustris sl	1		Calliergon giganteum	1	
fish bone	1		Carduus/Cirsium sp(p).	1	
fly puparia	1		Carex sp(p).	1	
Galeopsis Subgenus Galeopsis	1		Cerealina indet.	1	
Genista tinctoria (st fgts)	1		charcoal	1	max. size 15 mm
gravel	1	max. size 25 mm	Compositae (inv br)	1	
grit	1		Corylus avellana	1	
Homalothecium sericeum/ lutescens	1		Corylus avellana (ch)	1	
Humulus lupulus	1		Diphasium complanatum	1	
Hypnum cf. cupressiforme	1		earthworm egg caps	1	
Lapsana communis	1		eggshell fgts	1	max. size 2 mm
leather fgts	1	max. size 2 mm	Eleocharis palustris sl	1	
Leguminosae (pods/fgts)	1		fish bone	1	max. size 3 mm
Leucodon sciuroides	1		fish scale	1	
Linum usitatissimum			fly puparia	1	
(caps fgts)	1		Galeopsis Subgenus Galeopsis	1	
Lycopus europaeus	1		Galium sp(p).	1	
Malus sylvestris (endo)	1		Genista tinctoria (st fgts)	1	
mortar	1		glassy slag	1	max. size 2 mm
Neckera complanata	1		Gramineae	1	
Neckera crispa	1		gravel	1	max. size 20 mm
oyster shell fgts	1	max. size 5 mm	grit	1	
Polygonum aviculare agg.	1		herbaceous detritus	1	
Polygonum hydropiper	1		Hordeum sp(p).	1	
Polygonum lapathifolium	1		Humulus lupulus	1	
Polygonum persicaria	1		Hypnum cf. cupressiforme	1	
Potentilla cf. erecta	1		Lapsana communis	1	
pottery	1	max. size 15 mm	leather fgts	1	max. size 10 mm
Prunella vulgaris	1		Leguminosae (pods/fgts)	1	
Ranunculus flammula	1		marine mollusc shell fgts	1	max. size 5 mm
Ranunculus sardous	1		Marrubium vulgare	1	
Ranunculus Sect. Ranunculus	1		mortar	1	max. size 5 mm
Raphanus raphanistrum			Neckera complanata	1	
(pod segs/fgts)	1		oyster shell fgts	1	max. size 10 mm
Reseda luteola	1		Polygonum aviculare agg.	1	very small type
Rhytidadelphus sp(p).	1		Polygonum lapathifolium	1	
Rubia tinctorum	1		Polygonum persicaria	1	
Rumex acetosella agg.	1		pottery	1	
Rumex sp(p).	1		Prunella vulgaris	1	
Sambucus nigra	1		Ranunculus flammula	1	
Scorpidium scorpioides	1		Ranunculus Sect. Ranunculus	1	
Sonchus asper	1		Reseda lutea	1	
Stellaria media	1		Rubia tinctorum	1	
Thlaspi arvense	1		Rubus fruticosus agg.	1	
Triticum sp(p).	1		Sambucus nigra	1	
wood chips	1	max. size 15 mm	sand	1	
			Scorpidium scorpioides	1	
			Silene cf. alba	1	
			Spergula arvensis	1	
			Sphagnum sp(p). (lvs)	1	
			Stellaria media	1	
			Thuidium cf. tamariscinum	1	
			Urtica dioica	1	
			Urtica urens	1	
			Vicia faba	1	
<hr/>					
Context 5015, Sample 27/T (1.8 kg)					
wood fgts	3	max. size 35 mm			
Chenopodium album	2				
wood chips	2	max. size 35 mm			
Aethusa cynapium	1				
Agrostemma githago (sf)	1				

Context 5116, Sample 29/T

sand	3	
?human bone	1	
brick/tile	1	
Carex sp(p).	1	
charcoal	1	max. size 10 mm
marine mollusc shell fgts	1	
mortar	1	
Sambucus nigra	1	
tufa	1	
Urtica dioica	1	
wood fgts	1	very dec. fragments

Table 3. Complete list of invertebrate taxa recorded in samples from All Saints Church, York. Conventions: 'sp(?)' indicates 'probable additional taxon'; 'sp(?) indet.' indicates 'may be (or includes) previously listed taxon or taxa'. Order and nomenclature for Insecta follow Kloet and Hincks (1964-77). The ecological codes ascribed to species are appended. *—taxa not used in the calculation of main statistics.

CRUSTACEA - CLADOCERA		<i>Omalius</i> sp. indet.	rt
*Cladocera spp. F (ephippium)	oa-w	<i>Xylodromus concinnus</i> (Marsham)	rt-st
		<i>Coprophilus striatulus</i> (Fabricius)	rt-st
HEMIPTERA		<i>Carpelimus bilineatus</i> Stephens	rt-sf
Cimicidae sp.	oa-p	<i>Carpelimus fuliginosus</i> (Gravenhorst)	st
Saldidae sp.	oa-d	<i>Carpelimus pusillus</i> group	u
*Heteroptera sp. (nymph)	u	<i>Carpelimus</i> sp. indet.	u
<i>Conomelus anceps</i> (Germar)	oa-p	<i>Platystethus arenarius</i> (Fourcroy)	rf
*Coccoidea sp.	u	<i>Platystethus degener</i> Mulsant & Rey	oa-d
		<i>Platystethus nitens</i> (Sahlberg)	oa-d
LEPIDOPTERA		<i>Anotylus complanatus</i> (Erichson)	rt-sf
*Lepidoptera sp. (pupa)	u	<i>Anotylus nitidulus</i> (Gravenhorst)	rt-d
		<i>Anotylus rugosus</i> (Fabricius)	rt
DIPTERA		<i>Oxytelus sculptus</i> Gravenhorst	rt-st
*Syrphidae sp. (larva)	u	<i>Stenus</i> sp.	u
*?Melophagus ovinus (Linnaeus)	u	<i>Leptacinus</i> spp.	rt-st
*Melophagus ovinus (puparium)	u	<i>Gyrophypnus</i> sp.	rt
*Diptera sp. (adult)	u	Xantholininae sp.	u
*Diptera sp. (larva)	u	<i>Neobisnius</i> sp.	u
*Diptera sp. (pupa)	u	<i>Philonthus</i> spp.	u
*Diptera sp. (puparium)	u	Staphylininae spp.	u
		<i>Falagria</i> or <i>Cordalia</i> sp.	rt-sf
SIPHONAPTERA		? <i>Crataraea suturalis</i> (Mannerheim)	rt-st
* <i>Pulex irritans</i> Linnaeus	ss	Aleocharinae spp.	u
*Siphonaptera sp. indet.	u	Euplectini sp.	u
		Pselaphidae sp.	u
HYMENOPTERA		<i>Trox scaber</i> (Linnaeus)	rt-sf
*Proctotrupoidea sp.	u	<i>Aphodius granarius</i> (Linnaeus)	ob-rf
*Chalcidoidea sp.	u	<i>Aphodius</i> sp.	ob-rf
*Hymenoptera Parasitica sp.	u	<i>Cyphon</i> sp.	oa-d
* <i>Apis mellifera</i> Linnaeus	u	<i>Anobium punctatum</i> (Degeer)	l-sf
*Aculeata sp.	u	<i>Ptinus ?fur</i> (Linnaeus)	rd-sf
		<i>Lyctus linearis</i> (Goeze)	l-sf
COLEOPTERA		<i>Kateretes</i> sp.	oa-p-d
<i>Carabus nemoralis</i> Muller	oa	<i>Omosita discoidea</i> (Fabricius)	rt-sf
<i>Clivina ?fossor</i> (Linnaeus)	oa	<i>Cryptophagus scutellatus</i> Newman	rd-st
<i>Pterostichus ?melanarius</i> (Illiger)	ob	<i>Cryptophagus</i> spp.	rd-sf
? <i>Laemostenus terricola</i> (Herbst)	ss	<i>Atomaria</i> spp.	rd
<i>Helophorus aquaticus</i> or <i>grandis</i>	oa-w	<i>Orthoperus</i> sp.	rt
<i>Helophorus</i> sp.	oa-w	<i>Lathridius minutus</i> group	rd-st
<i>Cercyon analis</i> (Paykull)	rt-sf	<i>Corticaria</i> sp.	rt-sf
<i>Cercyon ?terminatus</i> (Marsham)	rf-st	<i>Typhaea stercorea</i> (Linnaeus)	rd-ss
<i>Cercyon ?unipunctatus</i> (Linnaeus)	rf-st	<i>Aglenus brunneus</i> (Gyllenhal)	rt-ss
<i>Cercyon</i> sp. indet.	u	<i>Tenebrio obscurus</i> Fabricius	rt-ss
<i>Cryptopleurum minutum</i> (Fabricius)	rf-st	<i>Phyllotreta nemorum</i> group	oa-p
<i>Acritus nigricornis</i> (Hoffmann)	rt-st	<i>Chaetocnema ?concinna</i> (Marsham)	oa-p
Histerinae sp.	rt	Halticinae sp.	oa-p
<i>Ochthebius</i> sp.	oa-w	<i>Apion</i> sp.	oa-p
<i>Catops</i> sp.	u	Curculionidae sp.	oa
<i>Omalius caesum</i> or <i>italicum</i>	rt-sf	*Coleoptera sp. (larva)	u
<i>Omalius ?rivulare</i> (Paykull)	rt-sf		

*Insecta sp. (larva) u

ARACHNIDA

*Aranae sp. u

*Acarina sp. u

Table 4. Main statistics for assemblages of adult beetles and bugs (excluding aphids and scale insects) from samples from All Saints Church, York, rounded to nearest whole figure. For explanation of abbreviations, see Table 5. Where alpha = 0 there were too few remains for a meaningful value to be calculated.

Context	4172	4172	5012	5012	5015	5015	Whole site
Sample	26	26	28	28	27	27	
Ext	/T(1)	/T(2)	/T(1)	/T(2)	/T(1)	/T(2)	
S	11	20	42	38	38	21	89
N	12	20	56	49	45	21	203
ALPHA	0	0	76	77	113	0	60
SEALPHA	0	0	22	25	45	0	7
SOB	2	6	9	9	7	3	23
PSOB	18	30	21	24	18	14	26
NOB	2	6	9	11	7	3	38
PNOB	17	30	16	22	16	14	19
ALPHAOB	0	0	0	0	0	0	25
SEALPHAOB	0	0	0	0	0	0	8
SW	0	1	2	0	1	1	3
PSW	0	5	5	0	3	5	3
NW	0	1	2	0	1	1	5
PNW	0	5	4	0	2	5	2
ALPHAW	0	0	0	0	0	0	0
SEALPHAW	0	0	0	0	0	0	0
SD	1	1	4	4	1	1	9
PSD	9	5	10	11	3	5	10
ND	1	1	6	6	2	1	17
PND	8	5	11	12	4	5	8
ALPHAD	0	0	0	0	0	0	0
SEALPHAD	0	0	0	0	0	0	0
SP	0	0	3	3	3	0	8
PSP	0	0	7	8	8	0	9
NP	0	0	3	3	3	0	9
PNP	0	0	5	6	7	0	4
ALPHAP	0	0	0	0	0	0	0
SEALPHAP	0	0	0	0	0	0	0
SM	0	0	0	0	0	0	0
PSM	0	0	0	0	0	0	0
NM	0	0	0	0	0	0	0
PNM	0	0	0	0	0	0	0
ALPHAM	0	0	0	0	0	0	0
SEALPHAM	0	0	0	0	0	0	0
SL	0	1	1	1	2	1	2
PSL	0	5	2	3	5	5	2
NL	0	1	1	1	2	1	6
PNL	0	5	2	2	4	5	3
ALPHAL	0	0	0	0	0	0	0
SEALPHAL	0	0	0	0	0	0	0
SRT	6	11	23	19	25	11	88
PSRT	55	55	55	50	66	52	99
NRT	7	11	34	27	32	11	122

Context	4172	4172	5012	5012	5015	5015	Whole site
Sample	26	26	28	28	27	27	
Ext	/T(1)	/T(2)	/T(1)	/T(2)	/T(1)	/T(2)	
PNRT	58	55	61	55	71	52	60
ALPHART	0	0	32	29	52	0	141
SEALPHART	0	0	11	12	21	0	27
SRD	1	4	7	5	5	0	22
PSRD	9	20	17	13	13	0	25
NRD	2	4	7	6	8	0	27
PNRD	17	20	13	12	18	0	13
ALPHARD	0	0	0	0	0	0	54
SEALPHARD	0	0	0	0	0	0	26
SRF	2	1	2	2	3	3	13
PSRF	18	5	5	5	8	14	15
NRF	2	1	2	3	3	3	14
PNRF	17	5	4	6	7	14	7
ALPHARF	0	0	0	0	0	0	0
SEALPHARF	0	0	0	0	0	0	0
SSA	1	10	17	13	21	12	36
PSSA	9	50	40	34	55	57	40
NSA	1	10	26	21	27	12	97
PNSA	8	50	46	43	60	57	48
ALPHASA	0	0	22	15	45	0	21
SEALPHASA	0	0	9	6	21	0	3
SSF	0	6	10	6	12	6	18
PSSF	0	30	24	16	32	29	20
NSF	0	6	13	12	17	6	54
PNSF	0	30	23	24	38	29	27
ALPHASF	0	0	0	0	0	0	10
SEALPHASF	0	0	0	0	0	0	2
SST	1	2	6	5	7	5	14
PSST	9	10	14	13	18	24	16
NST	1	2	12	7	8	5	35
PNST	8	10	21	14	18	24	17
ALPHAST	0	0	0	0	0	0	9
SEALPHAST	0	0	0	0	0	0	2
SSS	0	2	1	2	2	1	4
PSSS	0	10	2	5	5	5	4
NSS	0	2	1	2	2	1	8
PNSS	0	10	2	4	4	5	4
ALPHASS	0	0	0	0	0	0	0
SEALPHASS	0	0	0	0	0	0	0
SG	0	0	0	0	0	0	0
PSG	0	0	0	0	0	0	0
NG	0	0	0	0	0	0	0
PNG	0	0	0	0	0	0	0
ALPHAG	0	0	0	0	0	0	0
SEALPHAG	0	0	0	0	0	0	0

Table 5. Abbreviations for ecological codes and statistics used for interpretation of insect remains in text and tables. Lower case codes in parentheses are those assigned to taxa and used to calculate the group values (the codes in capitals). See Table 3 for codes assigned to taxa from All Saints Church, York. Indivs - individuals (based on MNI); No - number.

No taxa	S	Percentage of RT taxa	PSRT
Estimated number of indivs (MNI)	N	No RT indivs	NRT
Index of diversity (α)	alpha	Percentage of RT indivs	PNRT
Standard error of alpha	SE alpha	Index of diversity of RT component	alpha RT
No 'certain' outdoor taxa (oa)	SOA	Standard error	SEalphaRT
Percentage of 'certain' outdoor taxa	PSOA	No 'dry' decomposer taxa (rd)	SRD
No 'certain' outdoor indivs	NOA	Percentage of RD taxa	PSRD
Percentage of 'certain' outdoor indivs	PNOA	No RD indivs	NRD
No OA and probable outdoor taxa (oa+ob)	SOB	Percentage of RD indivs	PNRD
Percentage of OB taxa	PSOB	Index of diversity of the RD component	alphaRD
No OB indivs	NOB	Standard error	SEalphaRD
Percentage OB indivs	PNOB	No 'foul' decomposer taxa (rf)	SRF
Index of diversity of the OB component	alphaOB	Percentage of RF taxa	PSRF
Standard error	SEalphaOB	No RF indivs	NRF
No aquatic taxa (w)	SW	Percentage of RF indivs	PNRF
Percentage of aquatic taxa	PSW	Index of diversity of the RF component	alphaRF
No aquatic indivs	NW	Standard error	SEalphaRF
Percentage of W indivs	PNW	No synanthropic taxa (sf+st+ss)	SSA
Index of diversity of the W component	alphaW	Percentage of synanthropic taxa	PSSA
Standard error	SEalphaW	No synanthropic indivs	NSA
No damp ground/waterside taxa (d)	SD	Percentage of SA indivs	PNSA
Percentage D taxa	PSD	Index of diversity of SA component	ALPHASA
No damp D indivs	ND	Standard error	SEALPHASA
Percentage of D indivs	PND	No facultatively synanthropic taxa	SSF
Index of diversity of the D component	alphaD	Percentage of SF taxa	PSSF
Standard error	SEalphaD	No SF indivs	NSF
No strongly plant-associated taxa (p)	SP	Percentage of SF indivs	PNSF
Percentage of P taxa	PSP	Index of diversity of SF component	ALPHASF
No strongly P indivs	NP	Standard error	SEALPHASF
Percentage of P indivs	PNP	No typical synanthropic taxa	SST
Index of diversity of the P component	alphaP	Percentage of ST taxa	PSST
Standard error	SEalphaP	No ST indivs	NST
No heathland/moorland taxa (m)	SM	Percentage of ST indivs	PNST
Percentage of M taxa	PSM	Index of diversity of ST component	ALPHAST
No M indivs	NM	Standard error	SEALPHAST
Percentage of M indivs	PNM	No strongly synanthropic taxa	SSS
Index of diversity of the M component	alphaM	Percentage of SS taxa	PSSS
Standard error	SEalphaM	No SS indivs	NSS
No wood-associated taxa (l)	SL	Percentage of SS indivs	PNSS
Percentage of L taxa	PSL	Index of diversity of SS component	ALPHASS
No L indivs	NL	Standard error	SEALPHASS
Percentage of L indivs	PNL	No uncoded taxa (u)	SU
Index of diversity of the L component	alphaL	Percentage of uncoded indivs	PNU
Standard error	SEalphaL	No indivs of grain pests (g)	NG
No decomposer taxa (rt + rd + rf)	SRT	Percentage of indivs of grain pests	PNG

Table 6. Species lists in rank order for invertebrate macrofossils from samples from All Saints Church, York. For each sample assemblage the adult Hemiptera (bugs) and Coleoptera (beetles) are listed first, followed by the remaining invertebrates (*). Weight is in kilogrammes; n = minimum number of individuals; ec = ecological codes; sq = semi-quantitative (e = estimate; - = fully quantitative, m = 'many', translated as 15 individuals; s = several, translated as 6). For translation of ecological codes, see Table 5.

Context: 4172 Sample: 26/T(1) ReM: S
Weight: 1.00 E: 0.00 F: 0.00

Notes: One dish flot, few insects other than pupae. Recorded in flot and on filter paper. Preservation good to moderately good.

	n	sq	ec
Atomaria sp.	2	-	rd
Cercyon sp.	1	-	u
Catops sp.	1	-	u
Xylodromus ?concinus	1	-	rt-st
Platystethus arenarius	1	-	rf
Platystethus cornutus group	1	-	oa-d
Stenus sp.	1	-	u
Gyrophypnus sp.	1	-	rt
Staphylininae sp.	1	-	u
Aphodius granarius	1	-	ob-rf
Orthoperus sp.	1	-	rt
*Diptera sp. (pupa)	15	m	u
*Diptera sp. (puparium)	6	s	u
*Acarina sp.	1	-	u

Context: 4172 Sample: 26/T(2) ReM: S
Weight: 5.80 E: 0.00 F: 0.00

Notes: One dish flot, rather few insects. Recorded in flot and on filter paper; remains to tube. Fossils rather decayed and very fragmented. Some modern Nematocera adults

	n	sq	ec
Carabus nemoralis	1	-	oa
Pterostichus ?melanarius	1	-	ob
Ochthebius sp.	1	-	oa-w
Omalium caesum or italicum	1	-	rt-sf
Xylodromus ?concinus	1	-	rt-st
Carpelimus pusillus group	1	-	u
Platystethus nitens	1	-	oa-d
Anotylus complanatus	1	-	rt-sf
Stenus sp.	1	-	u
Staphylininae sp.	1	-	u
Falagria or Cordalia sp.	1	-	rt-sf
Aphodius sp.	1	-	ob-rf
Lyctus linearis	1	-	l-sf

Cryptophagus sp.	1	-	rd-sf
Atomaria sp.	1	-	rd
Lathridius minutus group	1	-	rd-st
Corticaria sp.	1	-	rt-sf
Typhaea stercorea	1	-	rd-ss
Aglenus brunneus	1	-	rt-ss
Curculionidae sp.	1	-	oa
*Diptera sp. (pupa)	15	m	u
*Acarina sp.	6	s	u
*Diptera sp. (puparium)	*6	s	u
*?Melophagus ovinus	1	-	u
*Siphonaptera sp.	1	-	u
*Diptera sp. (larva)	1	-	u
*Lepidoptera sp. (pupa)	1	-	u

Context: 5012 Sample: 28/T(1) ReM: S
Weight: 1.00 E: 0.00 F: 0.00

Notes: One dish flot; recorded in flot and on filter paper, remains to tube.

	n	sq	ec
Acritus nigricornis	4	-	rt-st
Xylodromus concinns	4	-	rt-st
Anotylus nitidulus	3	-	rt-d
Cercyon ?nalis	2	-	rt-sf
Carpelimus bilineatus	2	-	rt-sf
Carpelimus pusillus group	2	-	u
Anotylus complanatus	2	-	rt-sf
Neobisnius sp.	2	-	u
Philonthus sp. A	2	-	u
Conomelus anceps	1	-	oa-p
Clivina ?fossor	1	-	oa
Helophorus aquaticus or grandis	1	-	oa-w
Helophorus sp.	1	-	oa-w
Histerinae sp.	1	-	rt
Omalium caesum or italicum	1	-	rt-sf
Omalium sp.	1	-	rt
Coprophilus striatulus	1	-	rt-st
Carpelimus fuliginosus	1	-	st
Platystethus arenarius	1	-	rf
Platystethus ?degener	1	-	oa-d
Anotylus rugosus	1	-	rt
Philonthus sp. B	1	-	u

Staphylininae sp.	1	-	u	Anotylus rugosus	1	-	rt
Aleocharinae sp. A	1	-	u	Stenus sp.	1	-	u
Aleocharinae sp. B	1	-	u	Leptacinus sp. A	1	-	rt-st
Aleocharinae sp. C	1	-	u	Leptacinus sp. B	1	-	rt-st
Pselaphidae sp.	1	-	u	Gyrophypnus sp.	1	-	rt
Trox scaber	1	-	rt-sf	Neobisnius sp.	1	-	u
Aphodius sp.	1	-	ob-rf	?Philonthus sp.	1	-	u
Cyphon sp.	1	-	oa-d	Staphylininae sp. A	1	-	u
Anobium punctatum	1	-	l-sf	Staphylininae sp. B	1	-	u
Ptinus sp.	1	-	rd-sf	Aleocharinae sp. A	1	-	u
?Kateretes sp.	1	-	oa-p-d	Aleocharinae sp. B	1	-	u
Cryptophagus scutellatus	1	-	rd-st	Euplectini sp.	1	-	u
Cryptophagus sp. A	1	-	rd-sf	Aphodius sp.	1	-	ob-rf
Cryptophagus sp. B	1	-	rd-sf	Anobium punctatum	1	-	l-sf
Atomaria sp. A	1	-	rd	Kateretes sp.	1	-	oa-p-d
Atomaria sp. B	1	-	rd	Cryptophagus sp.	1	-	rd-sf
Lathridius minutus group	1	-	rd-st	Atomaria sp. A	1	-	rd
Corticaria sp.	1	-	rt-sf	Atomaria sp. B	1	-	rd
Aglenus brunneus	1	-	rt-ss	Lathridius minutus group	1	-	rd-st
Chaetocnema ?concinna	1	-	oa-p	Corticaria sp.	1	-	rt-sf
*Acarina sp.	6	s	u	Aglenus brunneus	1	-	rt-ss
*Diptera sp. (puparium)	6	s	u	Phyllotreta nemorum group	1	-	oa-p
*Coccoidea sp.	2	-	u	Apion sp.	1	-	oa-p
*Coleoptera sp. (larva)	2	-	u	Curculionidae sp.	1	-	oa
*Aculeata sp.	1	-	u	*Acarina sp.	15	m	u
*Chalcidoidea sp.	1	-	u	*Pulex irritans	1	-	ss
*Hymenoptera Parasitica sp.	1	-	u	*Aranae sp.	1	-	u
*Proctotrupoidea sp.	1	-	u	*Cladocera sp. F (ephippium)	1	-	oa-w
*Heteroptera sp. (nymph)	1	-	u	*Syrphidae sp. (larva)	1	-	u
*Diptera sp. (pupa)	1	-	u	*Diptera sp. (puparium)	1	-	u

Context: 5012 Sample: 28/T(2) ReM: S
Weight: 1.00 E: 0.00 F: 0.00

Context: 5015 Sample: 27/T(1) ReM: S
Weight: 1.00 E: 0.00 F: 0.00

Notes: One dish flot, preservation good. Recorded in flot and on filter paper, remains to tube. A modern mite noted. List includes some remains from residue ex ARH. Piece of ??weevil cuticle to own tube

Notes: Two-dish flot, preservation moderately good, though some fossils rotted. Recorded in flot and on filter paper, remains to tube

	n	sq	ec		n	sq	ec
Carpelimus ?bilineatus	4	-	rt-sf	Cryptophagus sp. B	3	-	rd-sf
Carpelimus fuliginosus	3	-	st	Cercyon ?nalis	2	-	rt-sf
Trox scaber	3	-	rt-sf	Carpelimus ?bilineatus	2	-	rt-sf
Platystethus degener	2	-	oa-d	Anotylus nitidulus	2	-	rt-d
Anotylus nitidulus	2	-	rt-d	Trox scaber	2	-	rt-sf
Aphodius granarius	2	-	ob-rf	Lathridius minutus group	2	-	rd-st
Ptinus ?fur	2	-	rd-sf	Cimicidae sp.	1	-	oa-p
Saldidae sp.	1	-	oa-d	Clivina ?fossor	1	-	oa
Clivina ?fossor	1	-	oa	Pterostichus ?melanarius	1	-	ob
?Laemostenus terricola	1	-	ss	Helophorus aquaticus or grandis	1	-	oa-w
Acritus nigricornis	1	-	rt-st	Cercyon analis	1	-	rt-sf
Histerinae sp.	1	-	rt	Cryptopleurum minutum	1	-	rf-st
Omalium sp.	1	-	rt	Xylodromus concinnus	1	-	rt-st
Carpelimus pusillus group	1	-	u	Coprophilus striatulus	1	-	rt-st
				Carpelimus fuliginosus	1	-	st

Carpelimus pusillus group	1	-	u	Carpelimus sp.	1	-	u
Platystethus arenarius	1	-	rf	Platystethus ?degener	1	-	oa-d
Anotylus complanatus	1	-	rt-sf	Oxytelus sculptus	1	-	rt-st
Anotylus rugosus	1	-	rt	Neobisnius sp.	1	-	u
Oxytelus sculptus	1	-	rt-st	Philonthus sp.	1	-	u
Gyrohypnus sp.	1	-	rt	Falagria or Cordalia sp.	1	-	rt-sf
Xantholininae sp.	1	-	u	Aleocharinae sp.	1	-	u
Neobisnius sp.	1	-	u	Euplectini sp.	1	-	u
?Cratarea suturalis	1	-	rt-st	Trox scaber	1	-	rt-sf
Aleocharinae sp.	1	-	u	Aphodius sp.	1	-	ob-rf
Aphodius sp.	1	-	ob-rf	Anobium punctatum	1	-	l-sf
Anobium punctatum	1	-	l-sf	Tenebrio obscurus	1	-	rt-ss
Ptinus sp.	1	-	rd-sf	*Pulex irritans	1	-	ss
Lyctus linearis	1	-	l-sf	*Acarina sp.	1	-	u
Omosita discoidea	1	-	rt-sf	*Cladocera sp. (ephippium)	1	-	oa
Cryptophagus sp. A	1	-	rd-sf				
Atomaria sp.	1	-	rd				
Orthoperus sp.	1	-	rt				
Corticaria sp.	1	-	rt-sf				
Aglenus brunneus	1	-	rt-ss				
Tenebrio obscurus	1	-	rt-ss				
Halticinae sp.	1	-	oa-p				
Apion sp.	1	-	oa-p				
*Acarina sp.	15	m	u				
*Diptera sp. (pupa)	15	m	u				
*Coccoidea sp.	6	s	u				
*Diptera sp. (puparium)	6	s	u				
*Hymenoptera Parasitica sp.	3	-	u				
*Proctotrupoidea sp.	3	-	u				
*Coleoptera sp. (larva)	2	-	u				
*Apis mellifera	1	-	u				
*Cladocera sp. F (ephippium)	1	-	oa-w				
*Diptera sp. (adult)	1	-	u				
*Siphonaptera sp.	1	-	u				
*Insecta sp. (larva)	1	-	u				
*Melophagus ovinus (puparium)	1	-	u				

Context: 5015 Sample: 27/T(2) ReM: S
 Weight: 0.80 E: 0.00 F: 0.00

Notes: One dish flot, preservation fairly good but some fossils rather fragmented. Recorded in flot and on filter paper, remains to tube

	n	sq	ec
Helophorus sp.	1	-	oa-w
Cercyon ?nalis	1	-	rt-sf
Cercyon ?terminatus	1	-	rf-st
Cercyon ?unipunctatus	1	-	rf-st
Omalium ?rivulare	1	-	rt-sf
Xylodromus concinnus	1	-	rt-st
Carpelimus ?bilineatus	1	-	rt-sf
Carpelimus fuliginosus	1	-	st
Carpelimus pusillus group	1	-	u